

1 Photo-electric effect


Photon's energy: $E = h\nu = hc/\lambda$

Photon's momentum: $p = h/\lambda = E/c$

Max. KE of ejected photo-electron: $K_{\max} = h\nu - \phi$

Threshold freq. in photo-electric effect: $\nu_0 = \phi/h$

Stopping potential: $V_o = \frac{hc}{e} \left(\frac{1}{\lambda}\right) - \frac{\phi}{e}$



de Broglie wavelength: $\lambda = h/p$

2 The Atom

Energy in n th Bohr's orbit:

$$E_n = -\frac{mZ^2e^4}{8\epsilon_0^2h^2n^2}, \quad E_n = -\frac{13.6Z^2}{n^2} \text{ eV}$$

Radius of the n th Bohr's orbit:

$$r_n = \frac{\epsilon_0h^2n^2}{\pi mZe^2}, \quad r_n = \frac{n^2a_0}{Z}, \quad a_0 = 0.529 \text{ \AA}$$

Quantization of the angular momentum: $l = \frac{nh}{2\pi}$


Photon energy in state transition: $E_2 - E_1 = h\nu$



Wavelength of emitted radiation: for a transition from n th to m th state:

$$\frac{1}{\lambda} = RZ^2 \left[\frac{1}{n^2} - \frac{1}{m^2} \right]$$

X-ray spectrum: $\lambda_{\min} = \frac{hc}{eV}$



Moseley's law: $\sqrt{\nu} = a(Z - b)$

X-ray diffraction: $2d \sin \theta = n\lambda$

Heisenberg uncertainty principle:


$$\Delta p \Delta x \geq h/(2\pi), \quad \Delta E \Delta t \geq h/(2\pi)$$

3 The Nucleus

Nuclear radius: $R = R_0A^{1/3}, \quad R_0 \approx 1.1 \times 10^{-15} \text{ m}$

Decay rate: $\frac{dN}{dt} = -\lambda N$

Population at time t : $N = N_0e^{-\lambda t}$



Half life: $t_{1/2} = 0.693/\lambda$

Average life: $t_{av} = 1/\lambda$

Population after n half lives: $N = N_0/2^n$

Mass defect: $\Delta m = [Zm_p + (A - Z)m_n] - M$

Binding energy: $B = [Zm_p + (A - Z)m_n - M]c^2$


Q-value: $Q = U_i - U_f$

Energy released in nuclear reaction: $\Delta E = \Delta mc^2$


where $\Delta m = m_{\text{reactants}} - m_{\text{products}}$.

4 Vacuum tubes and Semiconductors

Half Wave Rectifier:



Full Wave Rectifier:



Triode Valve:





Plate resistance of a triode: $r_p = \left. \frac{\Delta V_p}{\Delta i_p} \right|_{\Delta V_g=0}$

Transconductance of a triode: $g_m = \left. \frac{\Delta i_p}{\Delta V_g} \right|_{\Delta V_p=0}$

Amplification by a triode: $\mu = - \left. \frac{\Delta V_p}{\Delta V_g} \right|_{\Delta i_p=0}$

Relation between r_p , μ , and g_m : $\mu = r_p \times g_m$

Current in a transistor: $I_e = I_b + I_c$



α and β parameters of a transistor: $\alpha = \frac{I_c}{I_e}, \quad \beta = \frac{I_c}{I_b}$

Transconductance: $g_m = \frac{\Delta I_c}{\Delta V_{be}}$

Logic Gates:

		AND	OR	NAND	NOR	XOR
A	B	AB	A+B	\overline{AB}	$\overline{A+B}$	$A\overline{B} + \overline{A}B$
0	0	0	0	1	1	0
0	1	0	1	1	0	1
1	0	0	1	1	0	1
1	1	1	1	0	0	0

